VERSION 2

AUG 12, 2019

UC Davis - Massons Trichrome V.2

Jennifer
Rutkowski¹

¹University of California, Davis

Mouse Metabolic Phenotyping Centers
Tech. support email: info@mmpc.org

Lili Liang

ABSTRACT

Summary:

Massons trichrome staining is used for detection of collagen fibers in tissues such as skin, heart, etc. on formalin-fixed, paraffin-embedded sections, and may be used for frozen sections as well. The collagen fibers will be stained blue and the nuclei will be stained black and the background is stained red.

Modified from: Massons Trichrome UC Davis Clinical Pathology lab

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.56mg9c6

External link:
<https://mmpc.org/shared/document.aspx?id=259&docType=Protocol>

Protocol Citation: Jennifer Rutkowski 2019. UC Davis - Massons Trichrome.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.56mg9c6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Aug 05, 2019

Last Modified: Aug 12, 2019

Keywords: Massons
Trichrome

MATERIALS

MATERIALS

Weigerts Hematoxylin Contributed by
users

Bouins solution Contributed by
users

Biebrich Scarlet-Acid Fuchsin solution Contributed by
users

Phosphomolybdic- Phosphotungstic Acid solution Contributed by
users

Aniline Blue solution Contributed by
users

Acetic acid 1% Contributed by
users

coverslip Corning Catalog #2935-245

Reagent Preparation:

Reagent1: Biebrich Scarlet-Acid Fuchsin solution

Reagents and Materials:

Biebrich scarlet, aqueous, 1%	90 mL
Acid fuchsin, aqueous, 1%	10 mL
Glacial acetic acid	1.0 mL

Reagent 2: Phosphomolybdic- Phosphotungstic Acid solution

Reagents and Materials:

Phosphomolybdic acid	1.25 gm
Phosphotungstic acid	1.25 gm
Distilled water	50 mL

Procedure: Mix just before use

Reagent 3: Aniline Blue solution

Reagents and Materials:

Aniline blue	2.5 gm
Glacial Acetic acid	2.0 mL
Distilled water	100 mL

Procedure: Mix just before use

SAFETY WARNINGS

WARNING:

Formalin is, toxic, flammable and considered a carcinogen

All blood components and biological materials should be handled as potentially hazardous. Follow universal precautions established by CDC when handling and disposing of infectious agents.

- 1 Hydrate to water.
- 2 Place in Bouins for one hour in 55 degree oven or overnight at room temperature if needed.
- 3 Wash well in running water until yellow is gone.
- 4 Stain in Weigerts Hematoxylin for 10-20 minutes.
- 5 Wash well in running water for 10 minutes. Rinse distilled water.
- 6 Place slides in Biebrich Scarlet-Acid Fuchsin for 5 minutes.
- 7 Rinse in distilled water.

- 8 Place slides in the Phosphomolybdic-Phosphotungstic Acid Solution in a plastic coplin jar and put in the Bio Rad for 25 seconds using the total time setting. Let stand 30 seconds.
- 9 Place slides in Aniline Blue solution for 10 minutes.
- 10 Rinse in distilled water.
- 11 Place in 1% Acetic acid for 3 minutes if desired.
- 12 Dehydrate and coverslip.