VERSION 2

AUG 08, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.56vg9e6

External link:
<https://mmpc.org/shared/document.aspx?id=108&docType=Protocol>

Protocol Citation: John Rutledge 2019. UC Davis - Microvascular Permeability and Lipoprotein Flux. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.56vg9e6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Aug 05, 2019

Last Modified: Aug 08, 2019

PROTOCOL integer ID:
26549

UC Davis - Microvascular Permeability and Lipoprotein Flux V.2

John Rutledge¹

¹University of California, Davis

Mouse Metabolic Phenotyping Centers
Tech. support email: info@mmpc.org

Lili Liang

ABSTRACT

Summary:

One of the three indices of arterial function that are compromised to a varying degree in individuals with cardiovascular disease is vascular permeability. This assay measures vascular permeability (as flux of labeled large molecular weight molecules: i.e. albumin or dextran) and lipid permeability (as flux of labeled lipid) in coronary or carotid arteries.

Keywords: vascular permeability, lipid permeability, arterial function

MATERIALS

MATERIALS

☒ Krebs-Henseleit Solution **Contributed by users Catalog #See Below**

☒ FITC- Dextran **Sigma Aldrich Catalog #FD4, FD40S, or FD70**

☒ TRITC- Dextran **Sigma Aldrich Catalog #T1037 or T1162**

☒ FITC- Albumin **Sigma Aldrich Catalog #A9771**

☒ TRITC- Albumin **Sigma Aldrich Catalog #A2289**

☒ Alexa-546 label **Sigma Aldrich Catalog #10237**

☒ DiI labeled Lipid **Contributed by users Catalog #See protocol**

☒ pentobarbital **Cardinal Health**

☒ DMEM **Invitrogen - Thermo Fisher Catalog #11885**

☒ DPBS **Invitrogen - Thermo Fisher Catalog #14190**

☒ formaldehyde **Fisher Scientific Catalog #F79**

Reagent Preparation:

Reagent 1: 10 % formaldehyde

Formaldehyde (Fisher) is diluted to 10% in DPBS (Invitrogen)

Reagent 2: Krebs-Henseleit Solution

116 mM NaCl, 5 mM KCl, 2.4 mM $\text{CaCl}_2 \cdot \text{H}_2\text{O}$, 1.2 mM MgCl_2 , 1.2 mM NH_2PO_4 , and 11mM glucose

SAFETY WARNINGS

WARNING:

Formalin is, toxic, flammable and considered a carcinogen.

All blood components and biological materials should be handled as potentially hazardous. Follow universal precautions established by CDC when handling and disposing of infectious agents.

- 1 Mice are anesthetized with an intraperitoneal injection with 50 mg pentobarbital/kg weight.
- 2 All treatments are administered into the left femoral vein by bolus injection. FITC-albumin (40 mg/mL) in 100 μL :
 - a. PBS

- b. VLDL (150 mg/dL)
- c. VLDL (150mg/dL) + LpL (2 U/mL)
- d. LpL (2 U/mL) in PBS

- 3** Alternatively, the mouse is then infused at with 100 uL fluorescently labeled compound alone (FLC, see above) (40 mg/mL)
- 4** Excess FLC was removed from the vasculature by infusion with DMEM media for 20 min by infusion into the left ventricle of heart and followed by infusion of 10% formaldehyde for 20 min.
- 5** The microvascular rich tissues interest are immediately removed and fixed in 10% formaldehyde for two days.
 - a. microvascular tissues = brain, heart, and mesentery ect.
 - b. macrovascular tissues= common carotid arteries or aorta
- 6** The tissue is embedded in paraffin and sectioned to 5 μ m thickness.
- 7** Tissues sections are deparaffinized, rehydrated, and imaged using fluorescent microscopy.