

Aug 09, 2019 Version 2

UC Davis - Mouse Model Creation V.2

DOI

dx.doi.org/10.17504/protocols.io.56gg9bw



Kent Lloyd¹

¹University of California, Davis

Mouse Metabolic Phenotyping Centers
Tech. support email: info@mmpc.org



Lili Liang

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.56gg9bw

External link: <https://mmpc.org/shared/document.aspx?id=261&docType=Protocol>

Protocol Citation: Kent Lloyd 2019. UC Davis - Mouse Model Creation. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.56gg9bw>

Manuscript citation:

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: August 06, 2019

Last Modified: August 09, 2019

Protocol Integer ID: 26536

Keywords: Mouse Model



Abstract

Summary:

This is a generic protocol for making genetically-altered mouse models for research. Investigators who do not have a mutant mouse to send for testing at the MMPC @ UC Davis can ask for the mutant mouse to be derived by the MMPC @ UC Davis, through the UC Davis Mouse Biology Program. Genetically altered mice can be made on most genetic backgrounds, such as C57BL/6, FVB/N, B6D2F1 and other hybrids, and others. Transgenic (random genomic integration of expression-competent transgenes) and gene targeted (specific insertion and/or replacement of an endogenous genetic locus) mutant mice can be generated. "Knockout" mice can be made so as to delete endogenous gene expression ubiquitously and conditionally in either a specific cell or tissue type (spatial conditional) and/or at a specific age (temporal conditional). Numerous technologies are available for the derivation of mutant mice, including molecular construct design, gene targeted in embryonic stem (ES) cells, pronuclear microinjection of transgenic constructs into zygotes, ES cell microinjection into blastocysts, genome editing in ES cells and zygotes (e.g., TALENs, CRISPR/Cas9). Mutant mice can be genotype confirmed (see MMPC item D2004) and custom bred to generate cohorts of male and/or female mice for phenotyping at the MMPC @ UC Davis.



- 1 Depending on the type of mutant mouse requested, a number of protocols can be used. Investigators should consult with technical assistance staff at the MMPC @ UC Davis to develop their plan for deriving a mutant mouse.