MAY 17, 2019

UC Davis - Vertical Sleeve Gastrectomy

Kristin Evans¹

¹University of California, Davis

Mouse Metabolic Phenotyping Centers
Tech. support email: info@mmpc.org

Lili Liang

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.za3f2gn

External link:
<https://mmpc.org/shared/document.aspx?id=262&docType=Protocol>

Protocol Citation: Kristin Evans 2019. UC Davis - Vertical Sleeve Gastrectomy. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.za3f2gn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Mar 18, 2019

Last Modified: May 17, 2019

PROTOCOL integer ID:
21563

ABSTRACT

Summary:

Vertical sleeve Gastrectomy (VSG) is a common surgical treatment of obesity, type-2 diabetes, and other comorbidities of obesity. The VSG mouse surgical model provides researchers with a mechanism to evaluate mechanistic studies to explore the methods by which VSG bariatric surgery produces beneficial effects on obesity. The surgical procedure is performed under sterile conditions in accordance with AAALAC standards. Surgical animals will be able to ship 2 weeks after the post-operative monitoring period is completed. Cohorts of Vertical Sleeve gastrectomy (VSG), a pair fed SHAM, and an ad lib SHAM will be provided with each cohort ordered.

MATERIALS

MATERIALS

- ⊗ 0.9 % Sodium Chloride Injection USP B.Braun Medical Inc Catalog #NDC0264-4001-55
- ⊗ 4 x 4 non-woven gauze Veterinary Medical Central Services Catalog #LS2380
- ⊗ 2 x 2 non-woven gauze Veterinary Medical Central Services Catalog #LS2360
- ⊗ 20 cc syringe - Monoject Veterinary Medical Central Services Catalog #NS0440
- ⊗ 6-inch Non-sterile cotton swabs Fisher Scientific Catalog #23-400-101
- ⊗ 3-inch Non-sterile cotton swabs with pointed tip Fisher Scientific Catalog #23-400-118
- ⊗ 6-0 PDS suture noncutting needle (BV-1 taper double armed) Ethicon Catalog #Z117H

Keywords: Vertical sleeve
Gastrectomy (VSG)

7-0 PDS suture noncutting needle (BV-1 taper double armed) Ethicon Catalog #Z135H
Meloxicam SR (2 mg/ml) Zoopharm Catalog #Meloxicam SR (2 mg/ml)
Baytril (Enrofloxacin) 22.7 mg/ml Heartland Veterinary Catalog #2900-RX
70% isopropyl alcohol Contributed by users
Betadine surgical scrub Contributed by users
Betadine solution VWR international Ltd Catalog #AJ159701
Sterile drape General Econopack (STERIS Life Sciences) Catalog #88VCSTF
Clippers – Oster mini clipper Roboz Catalog #RS5903
Tissue glue Butler Schein (Henry Schein) Catalog #31477
Isoflurane USP Butler Schein (Henry Schein) Catalog #017579
Handheld cautery Roboz Catalog #RS300
Hemostats Roboz Catalog #RS7117(curved) RS7116(straight)
Micro-serrefines 18055-03 (straight)18055-05 (curved) Fine Science Tools Catalog #18055-03, 18055-05
Micro Needle Holder Fine Science Tools Catalog #12075-14
Spring Scissors Fine Science Tools Catalog #15024-10
Scissors Iris Fine Science Tools Catalog #14058-09
Dumont Forceps Fine Science Tools Catalog #11270-20
Micro Adson with fenestrated handle serrated Fine Science Tools Catalog #11020-12
Olsen-Hagar needle drivers Fine Science Tools Catalog #12002-14
Gavage needle Fine Science Tools Catalog #18061-75
Curved iris scissor Fine Science Tools Catalog #14079-10

⊗ Bead sterilizer (Germinator 500) Braintree Scientific Catalog #GER-5287-120V

⊗ Stereomicroscope Microscope Depot Catalog #GC01650

⊗ Surgisuite Multifunctional surgical platform Kent Scientific Catalog #Surgisuite

⊗ Isoflurane vaporizer VetEquip Catalog #Mobile Laboratory Animal Anesthe

Note:

B Braun Medical, Cite this, ([B Braun Sharing Expertise](#), [RRID:SCR_007148](#))
Fisher Scientific, [RRID:SCR_008452](#)

1 Preparation:

1. Animals should be fasted overnight to ensure that their stomach and intestines are as empty as possible.
2. Weight prior to surgery to calculate drugs (Meloxicam and Baytril dosing)
3. Autoclave all supplies and tools needed for the surgery, start the bead sterilizer for sterilizing instruments.
4. Prepare sterile field surrounding the microscope and surgisuite surgical platform. Turn on warming field.
5. Prepare sterilized supplies onto the sterile field that will be needed during the surgery.
6. Anesthetize mouse using isoflurane induction chamber 5% to effect. Once animal is unconscious they may be moved to the prep area and maintained under 1-1.5% isoflurane. Artificial tear salve is placed on each cornea to protect from drying.
7. Prepare the animal by clipping the hair from the surgical site (The length of the areas is from the knee to the armpit area and the width should cover the entire ventrum).
8. Perform a surgical scrub on the incision area using an initial wipe of Betadine surgical scrub followed by 70% alcohol working from the intended incision line outward. Continue preparing the surgical incision site by doing three successive scrub of Betadine solution followed by 70% alcohol using clean cotton tipped applicator for each application.
9. Administer Meloxicam SR 2 mg/kg subcutaneously in the rear dorsal area.
10. Spray the incision area using betadine solution before transferring the animal to the surgical area.

11. Surgeon will then wash and dry hands using soap and water and don sterile surgical gloves.

2 Surgery:

Stomach Isolation

1. Make small vertical midline skin incision from just below the xiphoid process the mid abdomen at the height of the top of the hip.
2. Cut along the linea alba from the xyphoid to just below the umbilical region
3. Gently using your gloved fingers or sterile cotton tip applicator, pull the intestine away to expose the stomach
4. Using a dry sterile cotton tip applicator gently place the applicator under the median and left liver lobes and under the stomach to elevate the stomach through the incision
5. Using a sterile cotton tip applicator bluntly dissect away the ligamentous fat from the stomach and tuck it back into the abdomen freeing all connections between the stomach and the liver
6. Remove the ligament to the liver that sits in the lesser curvature of the stomach using small iris scissors
7. Remove the ligament connecting the fundus to the body wall using small iris scissors
8. Place a cotton tip applicator under the fundus to elevate the vascular gastrosplenic ligament connection that attaches to the fundus.
9. Once the vascular connection is clearly exposed use the electrocautery to fuse the vascular connection and free the stomach

Vertical Sleeve Gastrectomy

1. Once the stomach is isolated, moisten the stomach and intestine with sterile saline using the gavage needle
2. Be careful not to pull the stomach out of the body cavity as this can injure the vagus nerves located here. Be careful of the pancreas attachment to the pyloric portion of the stomach. Avoid manipulation of this area.
3. The objective of the surgery is to remove 75-80% of the stomach
4. Place the isolated stomach on a 2 x 2 gauze and irrigate with sterile saline to prevent drying
5. Make an incision using the micro-scissors along the avascular seam of the greater curvature

of the stomach from the right of the esophagus to where the pancreatic vessels attach to the stomach.

6. Using a cotton tip applicator gently remove the stomach contents from the stomach incision onto the gauze

7. Replace the gauze and flush the stomach area with sterile saline

8. Replace the gauze with a fresh wet piece of gauze

9. Using sterile 7 French Polyethylene tubing, size the gastric pouch remnant from the esophagus to the pylorus

10. Using 7-0 PDS suture close the stomach at the edge of the sizing with the 6 French tubing using a simple continuous suture through both layers of stomach wall creating a diagonal line from the right of the esophagus to the connecting vessels of the pancreas. This leaves approximately 1/3 of the stomach cranial to the suture line.

11. Keep the remaining length of suture attached to close the stomach after removal of stomach tissue

12. Using micro scissors, cut away the portion of stomach tissue distal to the suture line being careful not to cut any of the previously laid suture material

13. Closure of the stomach is done using 7-0 PDS in a continuous mattress pattern moving from the pancreatic vessels to the esophagus.

14. Additional single interrupted suture are used if any gap areas are noted

15. The remaining stomach pouch should have a lumen diameter sized to 6 French and approximating 20-25% of the original stomach area

SHAM Vertical Sleeve Gastrectomy

1. Once the stomach is isolated, the stomach will have a gentle pressure applied using a pair of hemostats at a location that is approximately 2/3 cranial to the greater curvature.

2. The location of the clamp can be gauged by using a piece of 6 French tubing as a sizing indicator externally from the esophagus to the pylorus

3. Just below the gentle pressure line a two separate pieces of 7-0 suture will be run through the outer layer of stomach wall on both sides of the stomach

Surgical Closure

1. Perform an abdominal lavage with at least 25 ml of sterile saline using a gavage needle for at

least 4 gavages (total 100 ml)

2. Change the gauze under the stomach before doing the lavage and with each successive lavage to prevent stomach contents on the gauze from entering the abdomen
3. Place the stomach back into the abdomen
4. Suction remaining saline using sterile cotton tip applicators and gauze
5. Instill Baytril (20 mg/kg) diluted in sterile saline into the abdominal cavity before closing
6. Close the linea alba abdominal wall with 6-0 PDS in a simple continuous pattern
7. Close the skin with 6-0 PDS in a simple continuous pattern
8. Use tissue glue over the suture line

Post-operative care

1. Animals will be provided with slurry diet for 3 days post-operatively consisting of the powdered original diet or test diet indicated for the animals study combined with water (1:1 ratio). The slurry will be replaced daily.
2. Each animal will be observed once to twice daily for the first three days and then daily thereafter for 10 days (total of 14 of monitoring)
3. Animals will be assessed for pain level, incision status, status of eating, fecal production, and hydration status.
4. For the first three days following surgery subcutaneous saline (1 ml) will be administered to all surgical animals.
5. Animals will be transitioned onto regular chow pellets of the designated diet as indicated by the study at Day 3 post-surgery. From this point forward all animals will be pair feeding between CSG and pair fed Sham controls until shipping or study indicates
6. Animal body weights will be collected prior to surgery on the day of surgery, at 3 days post-operative, at 10 days post-operative, and at 14 days post-operative right before shipping or placement onto study.
7. All post-operative observation records, body weight records, and pair feeding information will be sent along with any shipped animal cohorts