VERSION 3

APR 12, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.z2jf8cn

External link:
<http://primerdigital.com/dna.html>

Protocol Citation: Ruslan Kalendar 2019. Universal DNA isolation protocol. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.z2jf8cn>

MANUSCRIPT CITATION:
 Kalendar R, Boronnikova S, Seppänen M. Isolation and Purification of DNA from Complicated Biological Samples. *Methods Mol Biol.* 2021;2222:57-67. doi: 10.1007/978-1-0716-0997-2_3. PMID: 33301087.

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Universal DNA isolation protocol V.3

Ruslan Kalendar¹

¹PrimerDigital

PrimerDigital

Ruslan Kalendar
PrimerDigital

ABSTRACT

The isolation of nucleic acids from a sample is an important step for many molecular biological applications and medical diagnostic assays. This protocol describes an efficient method for purification or/and isolation of nucleic acids from difficult animal tissues, plant material and other samples from which DNA extraction is generally regarded as being difficult which can contain impurities and inhibitors or interfering substances. This method is established acidic CTAB (with a pH value of 5 to 6.8) based extraction protocol that allows for reliable isolation of high molecular weight genomic DNA for removing contaminants from nucleic acids in a sample, e.g., environmental or biological samples such as soil, food, plant, animal or microorganism. DNA may quickly be extracted from samples without phenol, guanidine thiocyanate or 2-mercaptoethanol.

GUIDELINES

The procedure is suitable for all types of tissues from wide variety of animal, blood and plant species. All DNA extraction steps are performed at weak acid pH (HEPES free acid) and optionally with hot chloroform for 'difficult' samples, and at room temperature.

The following protocol is designed for small and large tissue samples (tissue volume 100-200 µl).

Note that isolating genomic DNA not requires gentle mixing because the DNA not be sheared by vortexing.

Protocol status: Working
We use this protocol and it's working

Created: Apr 12, 2019

Last Modified: Jul 08, 2022

PROTOCOL integer ID:
22315

Keywords: DNA extraction, CTAB, plant tissues, blood, herbarium specimens, difficult tissues

MATERIALS

MATERIALS

70% Ethanol Contributed by users

1L TE Buffer [10X] (Tris-EDTA) (100mM Tris base, 10mM EDTA, pH 8.0) G-Biosciences Catalog #786-034

CTAB (Hexadecyltrimethylammonium bromide) Bio Basic Inc. Catalog #CB0108.SIZE.500g

Chloroform:isoamyl alcohol (24:1) Bio Basic Inc. Catalog #CB0351.SIZE.200ml

HEPES, free acid Bio Basic Inc. Catalog #HB0264.SIZE.1Kg

Isopropanol Bio Basic Inc. Catalog #IB0918.SIZE.500ml

SAFETY WARNINGS

General rules for a laboratory safety should be followed.

BEFORE START INSTRUCTIONS

Required solutions:

- **CTAB** solution: 1.5% CTAB, 1.5 M NaCl, 10 mM Na₃EDTA, 0.1 M HEPES/MOPS (pH ~5.3);
100 ml: 1.5 g CTAB, 1.2 g HEPES-acid, 2 ml 0.5 M Na₃EDTA, 30 ml 5 M NaCl;
- Fresh 1xTE (1 mM EDTA, 10mM Tris-HCl, pH 8.0).

- 1 Eppendorf Safe-Lock microcentrifuge tube with tissue sample and glass ball (6 mm) freeze at -80°C, grind in the MM300 Mixer Mill for 2 min at 30 Hz. Alternatively, grind the sample in lysis solution.
- 2 In 2 ml tube with mechanically disrupted seeds/leaves/herbarium or DNA solution (CTAB purification) add 1 ml CTAB solution buffer with RNase A (the sample mass should not exceed 100 mg), vortex very well and incubate the samples at 60-65°C during 60-120 min or longer (long incubation increases DNA yield).

02:00:00

65 °C
- 3 Spin at maximum speed in a microcentrifuge for 2 minutes, transferred the upper aqueous layer to a new 2 ml microcentrifuge tube.

- 4 Transfer the entire clarified supernatant to a new 2 ml microcentrifuge tube which contains an equal volume of chloroform. Mix well for 3-5 minutes in the MM300 Mixer Mill at 30 Hz.
- 5 Spin at maximum speed in a microcentrifuge for 2 minutes.
- 6 Transfer the entire clarified upper aqueous layer to a new 2 ml microcentrifuge tube which contains an equal or half volume of 2-propanol and vortex thoroughly.
- 7 Centrifuge at maximum speed in a microcentrifuge for 2-5 minutes. A whitish DNA pellet should be visible.
- 8 Discard supernatant and wash the pellet by adding 1.8 ml 70% ethanol, vortex thoroughly. At this stage, DNA samples can be stored at room temperature or refrigerated.
- 9 Centrifuge at maximum speed for 2-5 min and carefully discard the supernatant by decanting or with a micropipette. A whitish DNA pellet should be visible during discarding a supernatant.
- 10 The DNA pellet does not dry and dissolved immediately in 300 µl 1xTE, pH 8.0 at 55°C for 10-20 minutes.