AUG 13, 2020

🌐 Universal sandwich enzyme linked immunosorbent assay for investigating protein-LG (SpLG) interactions with immunoglobulins using a SpA-HRP conjugate.

Angel A Justiz-Vaillant¹, Norma McFarlane-Anderson²

¹University of the West Indies St. Augustine;

²University of West Indies. Mona Campus

University of the West Indies

angel.vaillant@sta.uwi.edu

OPEN ACCESS

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

DOI:

dx.doi.org/10.17504/protocols.io.bjp4kmqw

Protocol Citation: Angel A Justiz-Vaillant, Norma McFarlane-Anderson 2020. Universal sandwich enzyme linked immunosorbent assay for investigating protein-LG (SpLG) interactions with immunoglobulins using a SpA-HRP conjugate.. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bjp4kmqw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Aug 13, 2020

Last Modified: Aug 13, 2020

PROTOCOL integer ID:
40412

1 This ELISA is used to study the interaction of a protein-LG (SpLG) with different immunoglobulin

preparations from mammalian species.

- 2 The 96 well microtitre plate is coated overnight at 4°C with 2 µg/µl per well of a mixture of protein-L and protein-G in carbonate-bicarbonate buffer pH 9.6.
- 3 Then plate is treated with bovine serum albumin solution and washed 4X with PBS-Tween.
- 4 50 µl of animal serum (1 mg/ml) is added and incubated for 1h at room temperature and the microplate is rewashed 4X with PBS-Tween.
- 5 Then 50 µl of peroxidase-labeled SpA conjugate diluted 1:5000 in PBS-non-fat milk is added to each well and incubated for 1h at RT. The plate is washed 4X with PBS-Tween.
- 6 50 µl of 4 mg/ml o-phenylenediamine solution (OPD) is added and the plate is incubated 15 minutes at RT in the dark.
- 7 The reaction is stopped with 50 µl of 3M H₂SO₄ solution.
- 8 The plate is visually assessed for the development of colour and read in a microplate reader at 492 nm.
- 9 A cut-off point should be calculated as the mean of the optical density of negative controls x 3. The higher the OD value the higher will be the affinity of SpLG to mammalian immunoglobulins.