

Dec 06, 2018 Version 1

Using fluorescent beads to measure feeding rate in *C. elegans* V.1

 PLOS Pathogens

DOI

[dx.doi.org/10.17504/protocols.io.v6ne9de](https://doi.org/10.17504/protocols.io.v6ne9de)

Emily Troemel¹

¹UCSD

 Emily Troemel

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.v6ne9de](https://doi.org/10.17504/protocols.io.v6ne9de)

External link: <https://doi.org/10.1371/journal.ppat.1007528>

Protocol Citation: Emily Troemel 2018. Using fluorescent beads to measure feeding rate in *C. elegans*. **protocols.io** <https://doi.org/10.17504/protocols.io.v6ne9de>

Manuscript citation:

Reddy KC, Dror T, Underwood RS, Osman GA, Elder CR, Desjardins CA, Cuomo CA, Barkoulas M, Troemel ER (2019) Antagonistic paralogs control a switch between growth and pathogen resistance in *C. elegans*. PLoS Pathog 15(1): e1007528. doi: [10.1371/journal.ppat.1007528](https://doi.org/10.1371/journal.ppat.1007528)

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: December 06, 2018

Last Modified: December 06, 2018

Protocol Integer ID: 18350

Attachments



DOCX

[Bead feeding assay.p...](#)

Attachments



DOCX

[Bead feeding](#)

[assay.p...](#)

0B

Attachments



DOCX

[Bead feeding](#)

[assay.p...](#)

0B

