MAY 16, 2019

🌐 Vandy – Energy Balance with Promethion

Louise Lantier¹

¹Vanderbilt University

Mouse Metabolic Phenotyping Centers

Tech. support email: info@mmpc.org

Lili Liang

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.yzxf7n

External link:

<https://mmpc.org/shared/document.aspx?id=241&docType=Protocol>

Protocol Citation: Louise Lantier 2019. Vandy – Energy Balance with Promethion.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.yzxf7n>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Mar 11, 2019

Last Modified: May 16, 2019

PROTOCOL integer ID:
21271

ABSTRACT

Summary:

The Promethion from Sable Systems (Las Vegas, NV) assesses several key metabolic characteristics of mice such as energy expenditure, preferred metabolic substrate, meal patterns, activity, live body mass. These are measured by individually housing the mice in the Promethion System cages for several days during which numerous parameters are continuously measured (food and water intake, weight, activity, O₂ and CO₂).

MATERIALS

MATERIALS

Promethion system **Sable Systems International**

Minispec Body Composition Analyzer **Bruker**

Scale **Contributed by users**

Keywords: Metabolic characteristics of mice, Promethion

- 1 One week prior to the experiment start date, mice are singly housed for acclimation.
- 2 On the day of the experiment start date, mice are weighed and body composition is assessed.
- 3 The mice are placed in the cages of the Promethion system, one mouse per cage. The cages are housed in a light and temperature controlled chamber. Default temperature is 23°C, but it can be set from 10 to 40°C to monitor mice in hypothermic or at thermoneutrality conditions. Likewise, the light cycle is set on a 12:12h cycle (6am-6pm) but can be modified to suit to study circadian rhythms.
- 4 Mice are left undisturbed for 5 days during which all the measurements are made.
- 5 After 5 days, the mice are removed from the cages, and body composition and weight are measured again.