MAY 16, 2019

Vandy - Hyperglycemic clamp

Li Kang¹

¹Vanderbilt University

Mouse Metabolic Phenotyping Centers

Tech. support email: info@mmpc.org

Lili Liang

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.yxefxje

External link:

<https://mmpc.org/shared/document.aspx?id=240&docType=Protocol>

Protocol Citation: Li Kang 2019. Vandy - Hyperglycemic clamp. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.yxefxje>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Mar 07, 2019

Last Modified: May 16, 2019

PROTOCOL integer ID:
21190

ABSTRACT

Summary:

Mice with catheters implanted in the jugular vein (infusions) and carotid artery (sampling) are used for this procedure. The hyperglycemic clamp is used to assess insulin secretory capacity in conscious mice in response to hyperglycemia. Plasma insulin and C-Peptide concentrations are measured at various times during the 2-hr clamp period.

Keywords: Hyperglycemic clamp

MATERIALS

MATERIALS

☒ Infusion Pumps Harvard Apparatus Catalog #PY8 70-2208

☒ Stand Fisher Scientific Catalog #14-670A

☒ Dual channel swivel Instech Solomon Catalog #375/D/22QM

☒ 3- and 4-way stainless steel connectors Ziggy's Tubes and Wires Catalog #HSC25 or HSC425

☒ Microrenthane tubing (0.033" OD) Braintree Scientific Catalog #MRE-033

☒ Glucose meter and strips ACCU-CHEK aviva

☒ Blunt needle with luer hub Ziggy's Tubes and Wires Catalog #LHN-E011041 25g x 0.5"

☒ Wire stainless steel Ziggy's Tubes and Wires Catalog #W020304V-1

☒ Clamp extension Fisher Scientific Catalog #05-769-7Q

☒ Connector hook Fisher Scientific Catalog #14-666-18Q

☒ 50% dextrose Contributed by users

Reagent Preparation:

Reagent 1: Donor Blood

1. Collect ~ 1 ml of blood from donor mouse in 0.5 ml EDTA tubes.
2. Centrifuge blood (1 min at 16,000 g) and save plasma for preparation of insulin (see below).
3. Resuspend red blood cells (RBC) with heparinized saline (10U/mL).
4. Centrifuge (1 min at 16,000 g), discard supernatant, and resuspend RBC with an equal volume of heparinized saline. Transfer resuspended RBC (donor blood) to a 1.5 ml tube

Note:

Fisher Scientific, [RRID:SCR_008452](https://doi.org/10.17504/protocols.io.yxefxje)

- 1 Surgical catheterization of the carotid artery and jugular vein in mice at least 5 days prior to the day of the study (refer to protocol for Surgical Catheterization of the Carotid Artery and Jugular Vein).
- 2 Weigh mouse and start fast (suggested starting time between 7:00 and 8:00 AM) by placing mouse in a plastic container with fresh bedding.

- 3** Mouse is hooked up to the swivel 4 hours into fasting and basal blood/plasma samples are collected 15 and 5 mins before the start of the clamp (refer to protocol for Hyperinsulinemic-Euglycemic Clamp for detailed set-up and connections).
- 4** After a total of 5 hr fast, variable infusion of 50% glucose starts. Arterial glucose is increased and maintained at 250-300 mg/dL.
- 5** Donor blood is infused to jugular vein catheter throughout the study to prevent a fall of hematocrit.
- 6** Plasma insulin and C-Peptide concentrations are measured at times as described in the following study sheet.
- 7** At the end of the study, mouse is anesthetized and tissues of interest are harvested and frozen in liquid nitrogen.

TIME (min)	SAMPLE (μ l)	GLUCOSE (mg/dl)	Time of infusion change	Glucose infusion rate		HCT	comments
				(μ l/min)	mg/kg/min		
-60	Place mouse in tub for acclimation						
-15	50 (G,I)						
-5	100(G,I,C)					*	Start donor blood
0	Variable Glucose Infusion (Clamp to ~300 mg/dl) Donor Blood :7ul/min						
5	50 (G,I)						
10	50 (G,I)						
15	100 (G,I,C)						
20	50 (G,I)						
30	5 (G)						
40	50 (G,I)						
50	5 (G)						
60	50 (G,I)						
70	5 (G)						
80	50 (G,I)						
90	5 (G)						
100	100 (G,I,C)						
110	5 (G)					*	
120	100 (G,I,C)						

I: sample for plasma insulin concentration (25 μ l plasma)

C: sample for C-peptide at t = -5,15,100, and 120 (50 μ l blood)