MAY 16, 2019

Vandy – IP glucose tolerance test

Louise Lantier¹

¹Vanderbilt University

Mouse Metabolic Phenotyping Centers
Tech. support email: info@mmpc.org

Lili Liang

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.yzzfx76

External link:
<https://mmpc.org/shared/document.aspx?id=238&docType=Protocol>

Protocol Citation: Louise Lantier 2019. Vandy – IP glucose tolerance test.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.yzzfx76>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Mar 11, 2019

Last Modified: May 16, 2019

PROTOCOL integer ID:
21273

ABSTRACT

Summary:

The glucose tolerance test measures the clearance of an intraperitoneally injected glucose load from the body. It is used to detect disturbances in glucose metabolism that can be linked to diabetes or metabolic syndrome. Animals are fasted for 5 hours, fasted blood glucose levels are determined before a solution of glucose is administered by intra-peritoneal (IP) injection. Subsequently, the blood glucose level is measured at different time points during the following 90 minutes.

MATERIALS

MATERIALS

⊗ 20% Dextrose **Contributed by users**

⊗ Glucometer and glucose strips **ACCU-CHEK aviva**

⊗ Syringes for IP injections **BD Biosciences**

⊗ Timer **Contributed by users**

⊗ Microvettes for blood collection **Sarstedt Catalog #16.444.100**

Note:

BD Biosciences [RRID:SCR_013311](#)

Keywords: Intraperitoneal
glucose tolerance test

- 1 Weigh the mice. For mice of differing fat mass, we recommend dosing the bolus of glucose on lean mass. This requires body composition.
- 2 Fast mice for 5 hours by transferring mice to individual cages or containers.
- 3 Prepare an experiment record sheet, sticks for glucose measurement and syringe for IP injection of the Dextrose solution.
- 4 Calculate and record the volume of 20% Dextrose solution required for each individual mouse for IP injection as follows:
 - a. To inject 2g of dextrose/kg body mass, the volume of the IP glucose injection is:
$$20\% \text{ Dextrose } (\mu\text{l}) = 10 \times \text{body weight (g)}$$
- 5 Cut the tip of the tail using clean surgical scissors. A small drop of blood (<5μl) is placed on the test strip of the blood glucose meter. This is the baseline glucose level (t = 0) and is recorded in the experiment record sheet.
- 6 Collect 40μL of whole blood in a tube for measurement of insulin. This is greatly facilitated by use of a collection tube such as Sarstedt's Microvette® CB 300 (cat # 16.443.100 for Heparin or cat # 16.444.100 for EDTA). Spin the blood and collect plasma in a clean tube (20μL). Store plasma on ice. Spin the blood 1min at 13k rpm and collect plasma in a new tube (25μL).
- 7 Inject the mouse intraperitoneally with the appropriate amount of glucose solution, as previously determined (point 4) and start the timer.
- 8 The blood glucose levels are measured at 5, 10, 15, 20, 30, 45, 60, 90 and 120 minutes after glucose gavage. Samples for insulin determination (20μl plasma) are collected at 10, 30, 60 and 120min (**Table 1**). For each sample, start the bleeding again by removing the clot from the first

incision, massage the tail if blood flow is inadequate. Place a small drop of blood on a new test strip. Results are recorded in the record sheet.

Mouse ID:			
Body Weight:			
Time (min)		Blood Glucose (mg/dL)	Comments
-5	G, I		
0	<i>Glucose injection (2g/kg body weight)</i>		
5	G		
10	G, I		
15	G		
20	G		
30	G, I		
45	G		
60	G, I		
90	G		
120	G, I		

G: glucose reading

I: Sample for Insulin (20µL plasma).

Table 1: IPGTT Record sheet

- 9 Ensure that further blood loss from the incision is minimal by briefly applying pressure to the incision after each measurement.

- 10 At the end of the experiment, mice are euthanized for tissue collection or placed in a clean cage with water and food available ad libitum for recovery. Monitor the mice carefully to observe any abnormal behavior. Plasma samples are stored in -20 or -80°C freezer until analysis.