

May 17, 2019

## Vandy – Post Clamp Anesthesia

DOI

[dx.doi.org/10.17504/protocols.io.zdnf25e](https://dx.doi.org/10.17504/protocols.io.zdnf25e)



Louise Lantier<sup>1</sup>

<sup>1</sup>Vanderbilt University

Mouse Metabolic Phenotyping Centers  
Tech. support email: [info@mmpc.org](mailto:info@mmpc.org)



Lili Liang

### Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.zdnf25e>

External link: <https://mmpc.org/shared/document.aspx?id=297&docType=Protocol>

**Protocol Citation:** Louise Lantier 2019. Vandy – Post Clamp Anesthesia. **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.zdnf25e>



**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**We use this protocol and it's working**

**Created:** March 21, 2019

**Last Modified:** May 17, 2019

**Protocol Integer ID:** 21646

**Keywords:** Post Clamp Anesthesia, post clamp anesthesia summary, anesthesia, pentobarbital, clamp, vandy, 35mg, mouse, important that the mouse

## Abstract

### Summary:

After the last sample is taken for the clamp, the mouse is anesthetized in order to harvest tissues. It is very important that the mouse remain alive during tissue harvesting for future signaling or metabolite analysis. The mouse is therefore given a sub-lethal dose of pentobarbital to induce anesthesia (~70mg/kg pi or 35mg/kg iv).

## Materials

### MATERIALS

⊗ Pentobarbital/ Nembutal® at 50mg/mL

⊗ Blunted needle

⊗ Sterile Saline solution

### Reagent Preparation:

#### **Reagent 1: Diluted Pentobarbital 5 mg/mL in saline from stock Pentobarbital (Nembutal®, 50mg/mL)**

1. For a 30mL Saline bottle: Inject 3.3mL of Stock Pentobarbital (50mg/ml), mix.
2. For a 10mL Saline bottle: Inject 1.1mL of Stock Pentobarbital, mix.

## Troubleshooting



- 1 Draw up the desired volume of **diluted Pentobarb (5mg/ml in saline)** in a blunted syringe.
  - To give 35mg/kg, multiply body weight (grams ) by 7, to get the volume to inject in uL.
  - Pentobarb (uL) = BW (g) \* 7
- 2 Steadily inject the pentobarbital in the jugular line of the catheterized mouse.
- 3 If the mouse is not completely anesthetized when taking the tissues, inject more by 50uL increments.

Concentration:		5 mg/ml						
Pentobarb mg/kg	BW mouse (g)							
	20	25	30	35	40	45	50	55
70	280	350	420	490	560	630	700	770
100	400	500	600	700	800	900	1000	1100