MAY 17, 2019

Vandy – Post Clamp Anesthesia

Louise Lantier¹

¹Vanderbilt University

Mouse Metabolic Phenotyping Centers
Tech. support email: info@mmpc.org

Lili Liang

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.zdnf25e

External link:
<https://mmpc.org/shared/document.aspx?id=297&docType=Protocol>

Protocol Citation: Louise Lantier 2019. Vandy – Post Clamp Anesthesia.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.zdnf25e>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Mar 20, 2019

Last Modified: May 17, 2019

PROTOCOL integer ID:
21646

ABSTRACT

Summary:

After the last sample is taken for the clamp, the mouse is anesthetized in order to harvest tissues. It is very important that the mouse remain alive during tissue harvesting for future signaling or metabolite analysis. The mouse is therefore given a sub-lethal dose of pentobarbital to induce anesthesia (~70mg/kg pi or 35mg/kg iv).

MATERIALS

MATERIALS

⊗ Pentobarbital/ Nembutal® at 50mg/mL **Contributed by users**

⊗ Blunted needle **Contributed by users**

⊗ Sterile Saline solution **Contributed by users**

Reagent Preparation:

Reagent 1: Diluted Pentobarbital 5 mg/mL in saline from stock Pentobarbital (Nembutal®, 50mg/mL)

1. For a 30mL Saline bottle: Inject 3.3mL of Stock Pentobarbital (50mg/ml), mix.
2. For a 10mL Saline bottle: Inject 1.1mL of Stock Pentobarbital, mix.

Keywords: Post Clamp
Anesthesia

- 1 Draw up the desired volume of **diluted Pentobarb (5mg/ml in saline)** in a blunted syringe.
 - To give 35mg/kg, multiply body weight (grams) by 7, to get the volume to inject in uL.
 - Pentobarb (uL) = BW (g) * 7
- 2 Steadily inject the pentobarbital in the jugular line of the catheterized mouse.
- 3 If the mouse is not completely anesthetized when taking the tissues, inject more by 50uL increments.

Concentration:		5 mg/ml						
Pentobarb mg/kg	BW mouse (g)							
	20	25	30	35	40	45	50	55
70	280	350	420	490	560	630	700	770
100	400	500	600	700	800	900	1000	1100