VERSION 2

OCT 18, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.8ezhtf6

External link:
<https://diacomp.org/shared/document.aspx?id=24&docType=Protocol>

Protocol Citation: Peter Mundel 2019. Vascular perfusion of mice.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.8ezhtf6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 18, 2019

Last Modified: Oct 18, 2019

PROTOCOL integer ID:
 28857

Vascular perfusion of mice V.2

Peter Mundel¹

¹Massachusetts General Hospital, Boston, MA

Diabetic Complications Consortium
 Tech. support email: rmcindoe@augusta.edu

Lili Liang

ABSTRACT

Summary:

This protocol describes the procedure for perfusion of mice prior to organ or tissue harvesting .

Diabetic Complications:

Cardiovascular

Retinopathy

Neuropathy

Nephropathy

Uropathy

Keywords: Vascular perfusion, cardiovascular, retinopathy, neuropathy, nephropathy, uropathy

MATERIALS

Reagents and Materials	Quantity Required
Perfusion Apparatus	1
Perfusion Bottles	2
Small diameter tubing (< 1cm)	
butterfly 23 gauge $\frac{3}{4}$	1
Ketamine	3 ml
Xylazine	1 ml
Plastic Tray	1
Phosphate Buffered Saline, pH 7.4	
PFA	
Sucrose	
70% alcohol	
Scissors	
Cotton swabs	

1 Preparation (see figure below for setup)

- ◆ Fix tubing (small diameter tubes, below 1 cm) to the bottles: 1 bottle for PFA, 1 bottle for sucrose
- ◆ Fill perfusion solutions in the bottles: solutions must be freshly prepared and filtered (using 0.22 μ m filter)
- ◆ Adjust perfusion pressure to 150-160 mm Hg
- ◆ Fix needle at the end of tubing after valve: butterfly 23 gauge $\frac{3}{4}$
- ◆ **Remove any air from all tubing!!**
- ◆ Anesthetic: ketamine-xylazine mix (150/10 mg/kg=0.1 ml/20 g BW)
- ◆ Mix 3 ml ketamine 100mg/ml + 1 ml xylazine 20mg/ml + 6 ml sterile saline
- ◆ Fix the anesthetized animal in a plastic tray
- ◆ Solution #1: 4 % PFA in PBS, pH 7.4
- ◆ Solution #2: 18% Sucrose in PBS, pH 7.4

2 Perfusion:

- ◆ Swab mouse with 70% alcohol to wet the fur, cut open abdominal skin by a longitudinal incision

- ◆ Remove skin and gut to expose abdominal aorta and Vena cava
- ◆ Clean the aort abdominalis and vena cava from connective tissue and fat with cotton swabs
- ◆ Clamp aorta abdominalis and vena cava **in the area of the iliac arteries**
- ◆ Insert the **butterfly 23 gauge 3/4** needle in the Aorta below the renal arteries
- ◆ Immediately cut open the Vena cava with small and sharp scissors: this should allow a rapid drain of the perfusion solutions!
- ◆ Watch the needle carefully during the entire perfusion!!!
- ◆ Start perfusion for 3 minutes with PFA, pH 7.4 at 37°C
- ◆ Switch valve to Solution #2:
- ◆ Without any interruption of pressure/flow perfuse for 5 minutes with sucrose, 7.4 at 37°C
- ◆ Stop perfusion, remove butterfly
- ◆ Cut out kidneys at renal hilum for further processing
- ◆ Before starting to perfuse the next animal rinse the tubing and needle extensively

3 Schematic of custom-made perfusion system