



Oct 03, 2024

Version 2

Vector Digestion and Purification V.2

DOI

<https://dx.doi.org/10.17504/protocols.io.261ge5dyog47/v2>

Carolina Lopez¹

¹Washington University



Cecilia Escudero

wustl

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.261ge5dyog47/v2>

Protocol Citation: Carolina Lopez 2024. Vector Digestion and Purification. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.261ge5dyog47/v2> Version created by **Carolina Lopez**

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: October 03, 2024



Last Modified: October 03, 2024

Protocol Integer ID: 109145

Keywords: purification protocol for plasmid digestion, plasmid digestion, purification protocol, vector digestion, purification

Abstract

Protocol for plasmid digestion and purification

Materials

Reagents:

- Restriction Enzymes (New England Biolabs)
- 10x Cutsmart Buffer (New England Biolabs)
- Agarose
- EtBr
- Promega Wizard SV Gel and PCR Purification Kit (A9282)



Isolate Digested Vector:

3h 2m

1 1. Digest Vector with New England Biolabs Restriction Enzymes:

3h 2m

Component	Volume (uL)
DNA Plasmid	X uL for 10 ug of Plasmid
10x Cutsmart Buffer	5
Enzyme 1	2.5
Enzyme 2	2.5
Nuclease Free H2O	40-X

2. Incubate for 02:00:00 at 37 °C .

3. Add 10 µL of 6x loading buffer to reaction and vortex briefly to mix.

4. Make 1% low melt-agarose gel.

a) Mix 1 g of Agar with 100 mL of TAE Buffer.

b) Microwave to boil agarose and let cool until you can touch bottle, but gel is not solid.

c) Add 1.5 µL of EtBr to agarose and pour into DNA gel mold with 10 well comb.

d) Let gel solidify.

5. Load 60 µL of reaction into well of gel

6. Run gel for 00:45:00 at 120V.

7. Visualize band with UV light and cut out section of gel with band with new razor blade and place in 1.5 mL tube.

8. Purify Band from gel with Promega Wizard SV Gel and PCR Purification Kit (A9282)

a) <https://www.promega.com/products/nucleic-acid-extraction/clean-up-and-concentration/wizard-sv-gel-and-pcr-clean-up-system/?catNum=A9281>

b) Weigh DNA gel fragment and add 10 µL of Membrane Binding Solution per 10 mg of gel slice.

c) Incubate mixture at 65 °C for 00:10:00 or until gel is completely melted.

d) Add melted gel mixture to SV minicolumn in Collection Tube and incubate at room temperature for 00:01:00 .

e) Centrifuge at max speed for 00:01:00 . Discard flowthrough and reinsert column into tube.

f) Add 700 µL of Membrane Wash Solution. Centrifuge at max speed for 00:01:00 . Discard flowthrough and reinsert column into tube.



g) Add  500 μ L of membrane wash solution. Centrifuge at max speed for

 00:01:00 . Discard flowthrough and reinsert column into tube.

h) Spin empty column for  00:01:00 at max speed to remove excess ethanol.

i) Transfer column to labelled  1.5 mL tube and add  35 μ L of NF H₂O.

Incubate for  00:01:00 at room temperature.

j) Centrifuge at max speed for  00:01:00 . Keep eluate and store at  -20 °C .