

Oct 06, 2019

## Virus titration

 [PLOS One](#)

DOI

[dx.doi.org/10.17504/protocols.io.7zhhp36](https://dx.doi.org/10.17504/protocols.io.7zhhp36)

Norfitriah Mohamed Sohaimi<sup>1</sup>, Mohd Hair Bejo<sup>1</sup>, Abdul Rahman Omar<sup>1</sup>, Aini Ideris<sup>1</sup>, Nurulfiza Mat Isa<sup>2</sup>

<sup>1</sup>Faculty of Veterinary Medicine, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.;

<sup>2</sup>Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.



Norfitriah Mohamed Sohaimi

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.7zhhp36](https://dx.doi.org/10.17504/protocols.io.7zhhp36)

External link: <https://doi.org/10.1371/journal.pone.0225863>

**Protocol Citation:** Norfitriah Mohamed Sohaimi, Mohd Hair Bejo, Abdul Rahman Omar, Aini Ideris, Nurulfiza Mat Isa 2019. Virus titration. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.7zhhp36>

**Manuscript citation:**

Sohaimi NM, Bejo MH, Omar AR, Ideris A, Isa NM (2019) Molecular characterization of fowl adenovirus isolate of Malaysia attenuated in chicken embryo liver cells and its pathogenicity and immunogenicity in chickens. PLoS ONE 14(12): e0225863. doi: [10.1371/journal.pone.0225863](https://doi.org/10.1371/journal.pone.0225863)

**License:** This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**We use this protocol and it's working**

**Created:** October 06, 2019

**Last Modified:** October 06, 2019

**Protocol Integer ID:** 28425



## Materials

### MATERIALS

✂ 96-well flat-bottom tissue culture plates with lids **Corning Catalog #3596**

Viral supernatant from passage 5 (CEL5), CEL10, CEL15, CEL20, CEL25, CEL30 and CEL35 were determined for titration by 50% tissue culture infective dose (TCID<sub>50</sub>) as described by Reed and Muench [1938]. Supernatant was diluted for 10 folds serial dilution and inoculated onto fresh CEL cells in 96 wells plate and monitored daily for CPE. Virus titre was calculated and expressed as TCID<sub>50</sub>/mL.

