VERSION 1

JUN 04, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.kfpctmn

External link:
<https://doi.org/10.1371/journal.pone.0192433>

Protocol Citation: João Vitor Molino 2018. Western blotting in *Chlamydomonas reinhardtii*. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.kfpctmn>

MANUSCRIPT CITATION:
 Molino JVD, Carvalho JCMd, Mayfield SP, Comparison of secretory signal peptides for heterologous protein expression in microalgae: Expanding the secretion portfolio for . PLoS ONE 13(2). doi: [10.1371/journal.pone.0192433](https://doi.org/10.1371/journal.pone.0192433)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Western blotting in *Chlamydomonas reinhardtii* V.1

PLOS One

João Vitor Molino¹

¹University of São Paulo

Joao Vitor Molino

Ronin Institute

ABSTRACT

This protocols describe the steps to perform a western blot in *Chlamydomonas reinhardtii* cell lysate and supernatant samples.

GUIDELINES

Prepare cell material by cultivating in liquid media.

BEFORE START INSTRUCTIONS

- Check antibody dilutions
- Check buffers disponibility
- Prepare 5% milk solution

Created: Oct 24, 2017

Last Modified: Jun 04, 2018

PROTOCOL integer ID:
8399

Sample preparation - Supernatant

1. Centrifuge algae culture at 2000xg for 10 min
2. Recover supernatant

00:10:00 Centrifugation time

Sample preparation - Lysate

1. Centrifuge algae culture at 2000xg for 10 min
2. Remove supernatant, and re-suspend cells in lysis buffer (50 mM Tris·HCL (pH 8.0), 0.1% Triton X-100), concentrating cells 100-fold.

00:10:00 Centrifugation time

Sonication

3. 1. Sonicate using appropriate sonication tip. Duty cycle: 0.5 | Amplitude: 20% | Duration: 30 s
2. Centrifuge for 15 min at 20000xg to remove cells debris and recover soluble proteins
3. Quantificate soluble protein

00:15:00 Centrifugation time

Gel electrophoresis | SDS-PAGE

4. 1. Load 30 µg of total soluble protein (TSP) per lane in a 12% SDS-PAGE
2. Transfer proteins to a nitrocellulose membrane
3. Block the membrane with 5% milk solution
4. Probe the desired protein with the specific antibody, diluted in 5% milk solution
5. Wash 2 times with TBST (0.2 M Tris, 1.37 M NaCl, 0.1% Tween-20, pH 7.6)
6. Add secondary antibody if required

30 µg total soluble protein per lane