NOV 21, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.8xehxje

External link:
<https://www.nature.com/articles/nprot.2015.122>

Protocol Citation: Jennifer B Treweek, Ken Y Chan, Nicholas C Flytzanis, Bin Yang, Benjamin E Deverman, Alon Greenbaum, Antti Lignell, Cheng Xiao, Long Cai, Mark S Ladinsky, Pamela J Bjorkman, Charless C Fowlkes, Viviana Gradinaru 2019. Whole-body tissue stabilization and selective extractions via tissue-hydrogel hybrids for high-resolution intact circuit mapping and phenotyping. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.8xehxje>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Nov 01, 2019

Last Modified: Nov 21, 2019

Whole-body tissue stabilization and selective extractions via tissue-hydrogel hybrids for high-resolution intact circuit mapping and phenotyping

Jennifer B Treweek¹, Ken Y Chan¹, Nicholas C Flytzanis¹, Bin Yang¹, Benjamin E Deverman², Alon Greenbaum¹, Antti Lignell³, Cheng Xiao¹, Long Cai³, Mark S Ladinsky¹, Pamela J Bjorkman¹, Charless C Fowlkes⁴, Viviana Gradinaru¹

¹Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, California, USA;

²Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, California, USA.;

³Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California, USA;

⁴Department of Computer Science, University of California, Irvine, California, USA

Optical Clearing of Tissue

Neurodegeneration Method Development Community

Viviana Gradinaru

ABSTRACT

To facilitate fine-scale phenotyping of whole specimens, we describe here a set of tissue fixation-embedding, detergent-clearing and staining protocols that can be used to transform excised organs and whole organisms into optically transparent samples within 1–2 weeks without compromising their cellular architecture or endogenous fluorescence. PACT (passive CLARITY technique) and PARS (perfusion-assisted agent release in situ) use tissue-hydrogel hybrids to stabilize tissue biomolecules during selective lipid extraction, resulting in enhanced clearing efficiency and sample integrity. Furthermore, the macromolecule permeability of PACT- and PARS-processed tissue hybrids supports the diffusion of immunolabels throughout intact tissue, whereas RIMS (refractive index matching solution) grants high-resolution imaging at depth by further reducing light scattering in cleared and uncleared samples alike. These methods are adaptable to difficult-to-image tissues, such as bone (PACT-deCAL), and to magnified single-cell visualization (ePACT). Together, these protocols and solutions enable phenotyping of subcellular components and tracing cellular connectivity in intact biological networks.

ATTACHMENTS

[nprot.2015.122.pdf](#)