VERSION 2

OCT 18, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.8e2htge

External link:
<https://diacomp.org/shared/document.aspx?id=63&docType=Protocol>

Protocol Citation: Frank Brosius 2019. WT-1 Staining Protocol for Podocytes.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.8e2htge>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 18, 2019

Last Modified: Oct 18, 2019

PROTOCOL integer ID:
28858

WT-1 Staining Protocol for Podocytes V.2

Frank Brosius¹

¹University of Arizona

Diabetic Complications Consortium
Tech. support email: rmcindoe@augusta.edu

Lili Liang

ABSTRACT

Summary:

This protocol describes a protocol used by some DiaComp members to detect podocyte nuclei in rodent glomeruli.

Edited by: Brosius Laboratory

Diabetic Complications:

Nephropathy

Keywords: podocyte nuclei in rodent glomeruli, nephropathy

MATERIALS

MATERIALS

- ☒ 75% 90% and 100% Ethanol **Contributed by users**
- ☒ PBS pH 7.4 **Contributed by users**
- ☒ Hydrogen peroxide **Contributed by users**
- ☒ Retrieve-All 1(1x) **Signet Pathology Systems Catalog #1912**
- ☒ 1% BSA in PBS **Contributed by users**
- ☒ WT1(C-19) rabbit polyclonal IgG **Santa Cruz Biotechnology Catalog #sc-192**
- ☒ DAB (Diaminobenzidine) Tablets **Sigma Aldrich Catalog #D-4293**
- ☒ Methanol **Contributed by users**
- ☒ Pemount **Contributed by users**
- ☒ Xylene **Contributed by users**
- ☒ Coverslips **Contributed by users**
- ☒ Staining Dish **Contributed by users**
- ☒ Staining Rack **Contributed by users**
- ☒ Staining Dish for boiling slides **VWR Scientific Catalog #25608-904**
- ☒ Vectastain ABC Kit Rabbit IgG **Vector Laboratories Catalog #PK-6101**

Reagent/Material	Quantity Required
Vectastain ABC Kit, Rabbit IgG	1
Retrieve-All 1(1x)	1
Staining Dish	12
Staining Rack	2

Note:

Santa Cruz Biotechnology ([RRID:SCR_008987](#))

Sigma-Aldrich ([RRID:SCR_008988](#))

Vector Laboratories Cat# PK-6101, [RRID:AB_2336820](#)

1 Day 1. Deparaffinize/hydrate

1. Dip in Xylene 5 minutes 2x (carry out in the hood)
2. Dip in 100% ETOH 5 minutes 2X
3. Dip in 95% ETOH 5 minutes once

4. Dip in 70% ETOH 5 minutes in once
5. Dip in dH₂O 5 minutes 2X
6. Dip in PBS 5 minutes once
7. Place the slides in jar with preheated Retrieve All 1 and incubate 2 hrs in 90°C water bath
8. Remove and let it cool down for 5-10 minutes
9. Pour off solution (can be reused 5-10 times) and add dH₂O
10. Rinse slides in two change of dH₂O 4-5 minutes each
11. Soak in PBS and store at 4°C.

2 Day 2. Immunoperoxidase

1. Blot dry slide (leave the section wet)
2. Incubate 20 minutes at room temperature (RT) with blocking serum (Follow directions of Vectastain ABS Kit for dilution).
3. Dilute WT-1 antibody (first antibody): 1:200 dilution in 1%BSA. Blot excess serum and incubate with first antibody for 2 hours at RT (2ug/section in 100ul).
4. Wash in PBS 5-10 minutes (2x changes) and blot dry (don't touch the section)
5. Incubate with second antibody for 1 hour (Follow directions of Vectastain ABS Kit for dilution)
6. Wash in PBS 5-10 minutes (2x changes)
7. Quench endogenous peroxidase by dipping in 1% H₂O₂ inmethanol for 45 minutes
8. Wash in PBS 10-15 minutes (2x changes)
9. Incubate with ABC Reagent for 1 hour (Follow direction ofVectastain ABS Kit for dilution).
10. Wash in PBS 5-10 minutes and prepare DAB Tablets (dissolve 1 set in 5 ml ddH₂O and filter it)
11. Staining: add 100ul or so DAB substrate to cover the whole section and check under the

microscope for podocyte nuclei (~1 minute).

12. Stop reaction in dH₂O

13. Dip in dH₂O for 5 minutes 2x

14. Dip in 70% ETOH 5 minutes once

15. Dip in 95% ETOH 5 minutes once

16. Dip in 100% ETOH 5 minutes 2X

17. Dip in Xylene 5 minutes 2x (carry out in the hood)

18. Mount the slides with permount.