OCT 12, 2019

Xylem inoculation with flg22

Cleo Bagchus¹

¹Wageningen University

iGEM Wageningen 2019

Cleo B.

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.752hq8e

Protocol Citation: Cleo Bagchus 2019. Xylem inoculation with flg22 . **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.752hq8e>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development
We are still developing and optimizing this protocol

Created: Oct 12, 2019

Last Modified: Oct 12, 2019

PROTOCOL integer ID:
28570

ABSTRACT

Protocol can be used after: Plant Infection with *Xanthomonas Campestris* Campestris

GUIDELINES

Depending on the country, *X. campestris* may be an quarantine organism. Please make sure you follow the specific guidelines of your country.

MATERIALS

sthree-week-old Brassica oleracea plants in climate chamber

Needle

1 uM flg22 at room temperature

Day 1

- 1 Place 15 µL of flg22 3 cm above the lesion on the central vein of the leaf.
- 2 With the needle puncture through the droplet into the central vein three times. Do not puncture the leaf completely. Due to a difference in pressure, the droplet should be sucked into the xylem.
- 3 Place a second 15 µL of flg22 3.5 cm above the lesion on the central vein of the leaf.
- 4 Repeat step 2.