

Oct 12, 2019

## Xylem inoculation with flg22

DOI

[dx.doi.org/10.17504/protocols.io.752hq8e](https://doi.org/10.17504/protocols.io.752hq8e)

Cleo Bagchus<sup>1</sup>

<sup>1</sup>Wageningen University

iGEM Wageningen 2019



Cleo B.

---

OPEN  ACCESS



**DOI:** [dx.doi.org/10.17504/protocols.io.752hq8e](https://doi.org/10.17504/protocols.io.752hq8e)

**Protocol Citation:** Cleo Bagchus 2019. Xylem inoculation with flg22 . **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.752hq8e>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** In development

We are still developing and optimizing this protocol

**Created:** October 12, 2019

**Last Modified:** October 12, 2019

**Protocol Integer ID:** 28570

### Abstract

Protocol can be used after: Plant Infection with *Xanthomonas Campestris* Campestris

## Guidelines

Depending on the country, *X. campestris* may be an quarantine organism. Please make sure you follow the specific guidelines of your country.

## Materials

sthree-week-old *Brassica oleracea* plants in climate chamber

Needle

1 uM flg22 at room temperature

## Day 1

- 1 Place  15 µL of flg22 3 cm above the lesion on the central vein of the leaf.
- 2 With the needle puncture through the droplet into the central vein three times. Do not puncture the leaf completely. Due to a difference in pressure, the droplet should be sucked into the xylem.
- 3 Place a second  15 µL of flg22 3.5 cm above the lesion on the central vein of the leaf.
- 4 Repeat step 2.