VERSION 2

JAN 01, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.vpfe5jn

External link:
<https://doi.org/10.1371/journal.pone.0209893>

Protocol Citation: Freddie-Jeanne Richard 2019. Y Choice test invertebrate.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.vpfe5jn>

MANUSCRIPT CITATION:
Beauché F, Richard F-J. The best timing of mate search in *Armadillidium vulgare* (Isopoda, Oniscidea). Plos One. 2013;8(3):e57737

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Nov 16, 2018

Y Choice test invertebrate V.2

PLOS One

Freddie-Jeanne Richard¹

¹Universtié de Poitiers

Freddie-Jeanne Richard

Universtié de Poitiers

ABSTRACT

Y choice test designed for terrestrial invertebrate

ATTACHMENTS

[Beauché Richard Plos one 2013.pdf](#)

Last Modified: Jan 01, 2019

PROTOCOL integer ID:
17863

Keywords: Behaviour , short distance, Perception

Device

- 1 Tests were performed using a Y-shaped choice chamber built in a plastic Petri dish (diameter: 8.7 cm; height: 1.2 cm; lid included) (Fig 2, from Beauché and Richard 2013 in Plos One). Rigid plastic tunnels were also used to create the device, and two plastic pipettes were sealed at the end of these tunnels to pulse air regularly into the system; the air was passed through sections (IIa) and (IIb) to spread the odor of the individuals.

Animals placement and monitoring

- 2 At the beginning of the experiment, the isopods were placed in the sections located in the three extreme parts of the tunnels: the two target females were placed in sections (IIa) and (IIb) (one in each section), and the tested male was placed in section (I). The male was then able to move into four other sections. The position of the target females in sections (IIa) and (IIb) was inverted between each replicate, and the target and tested individuals were used for only one test. The females were placed in their sections 15 minutes before the beginning of the test so that they could become accustomed to the new environment. The male was placed in section (I), and from the moment he entered the neutral section (NS), the time spent in the left section (LS) and the right section (RS) was monitored