



Jun 07, 2018

Yellow Fever Virus replication in cell culture

DOI

dx.doi.org/10.17504/protocols.io.pw6dphe

Izabela M Rezende¹, Livia Sacchetto

¹Stanford University School of Medicine

MRCA



Izabela M Rezende

Stanford University School of Medicine

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.pw6dphe

Protocol Citation: Izabela M Rezende, Livia Sacchetto 2018. Yellow Fever Virus replication in cell culture. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.pw6dphe>

Manuscript citation:

Persistence of Yellow fever virus outside the Amazon Basin, causing epidemics in Southeast Brazil, from 2016 to 2018. Rezende IM de*, Sacchetto L,* Mello ÉM de, Alves PA, Iani FC de M, Adelino TÉR, Duarte MM, Cury, ALF, Bernardes AFL, Santos TA, Pereira LS, Dutra MRT, Ramalho DB, Thoisy B, Kroon EG, Trindade GS, Drumond BP PLoS Neglected Tropical Disease 2018;12:e0006538. doi:10.1371/journal.pntd.0006538 *These authors contributed equally to this work

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: May 05, 2018

Last Modified: June 07, 2018

Protocol Integer ID: 11966



Abstract

In order to increase the number of viral particles a blind passage of sera might be performed in *Aedes albopictus* C6/36 cells.

Materials

MATERIALS

⊗ Fetal Bovine Serum, qualified **Life Technologies Catalog #10437-028**

⊗ PBS

⊗ Leibovitzs L-15 Medium **Merck MilliporeSigma (Sigma-Aldrich) Catalog #L4386**

⊗ Falcon® Serological Pipettes, 2 mL 1000 Pipettes **STEMCELL Technologies Inc. Catalog #38002**

⊗ Cell culture tubers **Techno Plastic Products (tpp) Catalog #91106**

⊗ C6/36 cells **ATCC**

⊗ 28°C incubator without CO2 **Thermo Fisher Scientific**

⊗ Invert Microscope

⊗ Falcon® Serological Pipettes, 5 mL **STEMCELL Technologies Inc.**

- 1 Remove and discard the cell culture media from the culture vessel.
- 2 Wash cells using PBS (approximately 2 mL per 10 cm² culture surface area). Gently add wash solution to the side of the vessel opposite the attached cell layer and wash the cells.
- 3 Shaking the flask slightly and using a pipette resuspend the cells in a minimal volume of pre-warmed Leibovit'z with 5% BFS and remove a sample for counting.
- 4 Using a cell counter, determine the number of cells/mL and adjust the number of cells to 65,000 C6/36 cells/mL.
- 5 Inoculate in the bottom of a cell culture tube 100,000 C6/36 cells in 1.5 mL of Leibovit'z medium supplemented with 5% BFS.
- 6 Keep the tubes inclined to 45 degrees at 28°C.
- 7 After 12-16 hours, remove and discard the media using a sterile pipette.
- 8 Wash the cells twice with phosphate buffered saline.
- 9 Add 100 µl of a solution containing 20 µL of sera and 80 µL of Leibovit'z medium.
- 10 Incubate the cells for 1 hour at 28°C.
- 11 After adsorption, add 1,5mL of Leibovit'z medium supplemented with 2% BFS and incubate the cells at at 28°C.
- 12 Observe the cells daily in an inverted microscope up to the observation of cytopathic effect or up to 10 days.



- 13 Collect the supernatant, make aliquots and kept them at -70°C . You can use this supernatant for other blind passages aiming viral isolation or to perform RNA extraction and further molecular analysis.