APR 08, 2019

Zebrafish Immunofluorescence Protocol

Kim Kobar¹

¹University of Wisconsin-Superior

Kim Kobar

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.zwif7ce

Protocol Citation: Kim Kobar
2019. Zebrafish
Immunofluorescence Protocol.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.zwif7ce>

License: This is an open
access protocol distributed
under the terms of
the [Creative Commons
Attribution License](#), which
permits unrestricted use,
distribution, and reproduction
in any medium, provided the
original author and source are
credited

Protocol status: Working
We use this protocol and it's
working

Created: Apr 08, 2019

Last Modified: Apr 08, 2019

PROTOCOL integer ID:
22186

Keywords: Zebrafish,
Immunofluorescence

MATERIALS

MATERIALS

☒ DMSO Fisher Scientific Catalog #BP231

☒ Goat Serum Gibco - Thermo Fischer Catalog #16210-064

☒ OmniPur BSA Fraction V (Bovine Serum Albumin) Heat Shock Isolation Contributed by users Catalog #2910-25GM

☒ Triton X-100 Fisher Scientific Catalog #BP151-100

☒ DPBS 10X with Calcium and Magnesium Gibco - Thermo Fischer Catalog #14-080-055

Recipes

IF Block solution

1. To make 10mL of blocking solution, combine and vortex the following in a 15mL conical:
2. 200µL goat serum (freezer)
3. 0.1g Bovine Serum Albumin (fridge)
4. 100µL DMSO (flammable cabinet)
5. 100µL 10% Triton X-100 (cabinet)
6. 1mL 10X PBS (cabinet)
7. 8.6mL DI H₂O
8. Solution should be cloudy, but if there are visible chunks floating around, place in 37°C water bath until uniform solution.
9. Store at 4° for 1 week or store at -20° for long term.

1X PBS

1. Add 9.6g of HyClone PBS (cabinet) to 1L of DI H₂O
2. Stir until dissolved and store at RT

PBS - 0.1% Triton

1. Add 100µL of Triton (cabinet) to 100mL 1X PBS

PBS - 2% Triton X-100

1. Add 2mL of Triton (cabinet) to 100mL 1X PBS

Day 1

- 1 Dechorionate embryos with two pairs of forceps.

Note

Under a dissecting microscope, pinch an embryo's chorion using a pair of forceps held in one of your hands. With the forceps in your other hand, pinch the chorion near to the original pinch and gently tear the chorion by separating your hands. Repeat pinching and tearing chorions with forceps until the embryos are dechorionated.

- 2 Using a large bore pipette, transfer dechorionated embryos into labelled microcentrifuge tubes, up to 50 embryos/tube.

Note

The labels should include: # of embryos, age in hpf, what was injected into them & how much, and the date the embryos were injected. Label the top and side of the tube.

- 3 Remove as much liquid as possible from the microcentrifuge tubes with a small bore pipette.
- 4 Fix dechorionated embryos in ~1mL of 4% PFA (if used every week - stored in 4°C; new aliquots are stored long term at -20°C for 1 hour at RT on the rocker). Tubes should be placed on their side whenever they are rocking. Orient the tubes perpendicular to the rocking motion such that the embryos rock from side to side in the tube rather than from cap to bottom.

Safety information

Wear gloves when working with PFA

- 5 Discard PFA from tubes into PFA waste using a small bore pipette. Take care to not pipet or damage any embryos when removing solutions.
- 6 Rinse in ~1mL PBS 2 x 5 mins at RT on the rocker. Remove solution with a small bore pipette each time before adding new solution.

00:05:00 First Wash

00:05:00 Second Wash

Note

Embryos can then be stored in fresh PBS for 2 weeks max at 4°C (in the box labeled "Fixed Embryos" - be sure to add each tube to the "Fixed Embryos Log."

START HERE when using previously fixed embryos:

7 For 24-48 hpf embryos:

- wash in ~1mL DI H₂O for 5 min at RT on the rocker.

For ~72 hpf embryos:

- wash 5 min in DI H₂O at RT on the rocker
- put in 100% acetone at -20°C for 5 min,
- then wash in DI H₂O again for 5 min at RT on the rocker.

8 Incubate for 1 hour in 500 ul IF Block solution at RT on the rocker.

Note

IF Block is stored in 15 ml conicals in the -20°C - however, keep IF Block at 4°C from now until done making the 2° Antibody Solution on Day 2 (this is to avoid repeated freeze/thaw cycles).

9 Remove IF Block solution from embryos and discard in waste beaker.

10 Replace with 1° Antibody solution: 100 ul per tube of embryos (1° antibody is diluted in IF Block solution, the amount of dilution varies with each antibody - see the table at the end of this protocol to determine correct concentration).

11 Incubate overnight at 4 °C

Day 2

12 Remove the 1° Antibody Solution and discard in waste beaker.

13 Rinse all day in 100µL-500µL PBS- 0.1% Triton at room temperature on the rocker.

Note

Change the solution 3-4 times throughout the day, ensuring to remove solution before adding new solution.

14 Remove PBS-0.1% Triton and replace with 2° Antibody Solution.

Note

Choose the Dylight488 (green) conjugated 2° antibody made against the species in which your primary antibody was produced. **However, if doing a dual stain, choose Dylight 594 (red).** Secondary antibodies are diluted at 1:750 in IF block (see table): 100 ul per tube of embryos.

15 Incubate **in dark** (wrap with tin foil) overnight at 4°C.

Note

Keep in dark as much as possible from now on out.

Day 3

16 Remove solution in tubes with small bore pipette. Rinse in PBS- 0.1% Triton 2 x 5min at RT on the rocker.

17 Store in at least 200 ul PBS at 4°C.

Note

Try to image within a week to avoid degradation.

Antibody Dilution Table

- 18** Primary Antibody Dilutions: prepare ~100 ul per tube of 50 embryos
Most primary antibodies are stored in the "Research S18" box in 4C fridge.

Example: $\frac{1}{20} = \frac{5 \text{ ul antibody}}{95 \text{ ul IF Block}}$

Name	Dilution	Species	Vendor/Product #
F310	1:20/ 1:50	mouse	
A4.10 25	1:20	mouse	
Phalloidin (Actin)	1:20		
PH3	1:500	rabbit	
CC3	1:500	rabbit	
Dylight 488 α -rabbit	1:750		
Dylight 488 α -mouse	1:750		
Dylight 594 α -rabbit	1:750		
Dylight 594 α -mouse	1:750		
HRP α -rabbit

HRP α- mous e			
------------------------	--	--	--